arXiv:q-alg/9505017v1 17 May 1995

LMU–TPW 95–6

$ISO_q(3)$ and $ISO_q(2,1)$

Sebastian Sachse, Ralf Weixler
Sektion Physik der Universität München
Lehrstuhl Professor Wess, Theresienstraße 37
D-81245 München , Germany

February 9, 2008

Abstract

We prove the embedding of $ISO_q(3) \hookrightarrow ISU^{ex}_{\sqrt{q}}(2)$ and $ISO_q(2,1) \hookrightarrow ISL^{ex}_q(2,R)$ as *-algebras and give a Hilbert space representation of $ISU^{ex}_{\sqrt{q}}(2)$.

1 Introduction

The inhomogenized extensions of a large list of standard quantizied Lie groups [1] have been given in [2, 3, 4, 5, 7, 8]. They form quantizied versions of the classical inhomogeneous groups. For a real deformation parameter q the representation theory of the homogeneous parts (e.g. corepresentations of the function algebra) is basically the same as for the classical groups, whereas for q root of unity it is completely different. The representation theory of the noncommutative function algebra however differs for any $q \neq 1$ from the classical situation. Its relevance stems from the question whether a deformation exists on the C^*-algebra level. [6]

In part 2 we recall the properties of inhomogeneous quantum groups. In the 3rd. part we examine the algebraic embedding of the $ISO_q(3)$ into $ISU^{ex}_{\sqrt{q}}(2)$ and

$ISO_q(2,1)$ into $ISL_q^{ex}(2,R)$. Here the "extended inhomogeneous" quantum algebra IG^{ex} designates inhomogeneous quantum algebras containing two sets of coordinate functions.

In the last chapter we examine the representation theory of the $ISU_{\sqrt{q}}^{ex}(2)$.

2 The Hopf algebra structure of inhomogeneous quantum groups

Quantum groups may be considered to be deformations of the function algebra over the corresponding Lie groups. The deformation is given by a parameter $q \epsilon \mathbb{C}$ which has to be further restricted in order to get special cases of deformations. Quantum groups exhibit a Hopf algebra structure. The noncommutative algebra structure is controlled by an $\hat{R}$–matrix fulfilling the Quantum Yang–Baxter equation $\hat{R}_{12}\hat{R}_{23}\hat{R}_{12} = \hat{R}_{23}\hat{R}_{12}\hat{R}_{23}$. In this paper we refer to $\hat{R}$–matrices in their standard form given in [1]. They are defined by their projector decomposition making use of the antisymmetrizer $\hat{A}_q{}^{ij}{}_{kl}$, the symmetrizer $\hat{S}_q{}^{ij}{}_{kl}$ and the trace projector $\hat{T}_q{}^{ij}{}_{kl} \propto C^{ij}C_{kl}$ with the metric C_{ij}, existing for the q–orthogonal groups only.

$$\hat{R}_q{}^{ij}{}_{kl} = \begin{cases} q\hat{S}_q{}^{ij}{}_{kl} - q^{-1}\hat{A}_q{}^{ij}{}_{kl} & (\text{ for } SL_q(N)) \\ q\hat{S}_q{}^{ij}{}_{kl} - q^{-1}\hat{A}_q{}^{ij}{}_{kl} + q^{1-N}\hat{T}_q{}^{ij}{}_{kl} & (\text{ for } SO_q(N)) \end{cases} \tag{1}$$

The algebra relations for the generators $M^i{}_j$ of the unital $\mathbb{C}$ -algebra $\mathcal{A}$ are:

$$\hat{R}_q{}^{ij}{}_{j'i'} M^{j'}{}_{j''} M^{i'}{}_{i''} = M^i{}_{i'} M^j{}_{j'} \hat{R}_q{}^{i'j'}{}_{j''i''} \tag{2}$$

and

$$\begin{cases} \det M = \frac{(-1)^{N-1}}{[N]!}\epsilon^{k_1 \ldots k_N} M^{l_1}{}_{k_1} \ldots M^{l_N}{}_{k_N} \epsilon_{l_1 \ldots l_N} = \mathbf{1} & (SL_q(N)) \\ C_{ij} M^i{}_{i'} M^j{}_{j'} = C_{i'j'}\mathbf{1} & (SO_q(N)) \end{cases} \tag{3}$$

For the unimodularity condition we use the q–antisymmetric tensors ϵ_q defined in [10].

In order to obtain inhomogeneous quantum groups the set of generators has to be enlarged not only by the coordinate functions x_i but by an invertible scaling operator $\bar{w}$ as well. Its existence is required by consistency of the comultiplication. The additional algebra relations of the extended Hopf algebra $\mathcal{A}^I$ are:

$$\begin{array}{llll} (i) & x^i M^j{}_k = \gamma \hat{R}_q{}^{ij}{}_{lm} M^l{}_k x^m & (iv) & w M^i{}_j = M^i{}_j w \\ (ii) & \bar{w} w = \mathbf{1} & (v) & \bar{w} x^i = \dfrac{q}{\gamma} x^i \bar{w} \\ (iii) & \bar{w} M^i{}_j = M^i{}_j \bar{w} & (vi) & w x^i = \dfrac{\gamma}{q} x^i w \end{array} \tag{4}$$

with

$$\gamma = \begin{cases} q^{-1/N} & (\text{ for } SL_q(N)) \\ 1 & (\text{ for } SO_q(N)) \end{cases} \tag{5}$$

The comultiplication $\Phi : \mathcal{A}^I \to \mathcal{A}^I \otimes \mathcal{A}^I$, counit $e : \mathcal{A}^I \to \mathbb{C}$ and the antipode $\kappa : \mathcal{A}^I \to \mathcal{A}^I$ are very easily given in matrix notation.

With

$$M^I = \begin{pmatrix} \bar{w}M & \mathbf{x} \\ 0 & 1 \end{pmatrix} \tag{6}$$

we get

$$\Phi(M^I) = M^I \dot{\otimes} M^I \tag{7}$$

and

$$e(M^I) = \begin{pmatrix} E & \mathbf{0} \\ 0 & 1 \end{pmatrix} \tag{8}$$

with the unity matrix E.

The antipode is given as

$$\kappa(M^I) = \begin{pmatrix} \kappa(M)\bar{w} & -\kappa(M)\bar{w}\vec{x} \\ 0 & \mathbf{1} \end{pmatrix}. \tag{9}$$

2.1 Complex conjugation

The $*$-operations on $\mathcal{A}$ are defined quite differently in the cases $q\epsilon\mathbf{R}^+$ and $|q| = 1$.
a) $q\epsilon\ \mathbf{R}^+$

With the unitarity condition $(M^i{}_j)^* = \kappa(M^j{}_i)$, the quantum group $SL_q(N)$ becomes a $SU_q(N)$ quantum group. The same *-structure holds for the orthogonal quantum groups $SO_q(N)$.

b) $|q| = 1$

For such q the R_q matrix has the property $R_q^* = R_q^{-1}$. With the reality condition $(M^i{}_j)^* = M^i{}_j$ one finds the real representation of the quantum group $SL_q(N)$ called $SL_q(N, R)$ and the orthogonal quantum groups in this case have a metric which is indefinite, i.e. for N even we get $SO_q(n, n)$ and for N odd $SO_q(n, n+1)$, with $N = 2n$ or $N = 2n+1$ respectivly.

The complex conjugation for the inhomogeneous extensions of these function algebras have to be treated separatly as well.

a) In the case $q\epsilon\ \mathbf{R}^+$ we have to enlarge the generating set of the *-Hopf algebra $\mathcal{A}^I$ by the conjugate coordinate functions $\bar{x}_i$. The additional algebra relations are:

$$\begin{array}{llll} (i) & M^l{}_s\bar{x}_j = \dfrac{1}{\gamma}\hat{R}_q{}^{al}{}_{ij}\bar{x}_a M^i{}_s & (iii) & \bar{w}\bar{x}_i = \dfrac{q}{\gamma}\bar{x}_i\bar{w} \\ (ii) & w\bar{x}_i = \dfrac{\gamma}{q}\bar{x}_i w & (iv) & x^i\bar{x}_j = \dfrac{1}{q}\bar{x}_a x^b \hat{R}_q{}^{ai}{}_{bj}. \end{array} \tag{10}$$

The comultiplication of $\bar{x}_i$ follows from being an *-homomorphism and the antipode from the fact that $\kappa \circ * \circ \kappa \circ * = \mathrm{id}$.

b) When $|q| = 1$ the coordinate functions may be chosen to be real ($x_i^* = x_i$), since applying the * to (4(i)) and taking into account that $R_q{}^{ij*}_{kl} = R_q^{-1ij}{}_{kl}$, we get:

$$M^j{}_k x^i = \gamma^{-1} R_q^{-1ji}{}_{lm} x^m M^l{}_k \tag{11}$$

or

$$\gamma \hat{R}_q{}^{rs}{}_{ji} M^j{}_k x^i = \delta^r_m \delta^s_l x^m M^l{}_k. \tag{12}$$

3 Algebraic embedding

The $\hat{R}-$matrix of $SU_{\sqrt{q}}(2)$ is decomposed using the q-antisymmetric ε-tensor:

$$\hat{R}^{\mu\nu}_{\sqrt{q}\rho\sigma} = \sqrt{q}\delta^{\mu}_{\rho}\delta^{\sigma}_{\nu} + \varepsilon^{\mu\nu}\varepsilon_{\rho\sigma}, \tag{13}$$

with

$$\epsilon_{\mu\nu} := \begin{pmatrix} 0 & q^{-1/4} \\ -q^{1/4} & 0 \end{pmatrix}_{\mu\nu} = -\epsilon^{\mu\nu}$$

For the homogeneous parts the embedding $SO_q(3) \hookrightarrow SU_{\sqrt{q}}$ is well known [9]. With the q-deformed Clebsch-Gordan-coefficients $c^i_{\mu\nu}$ of the product decomposition of the $SU_{\sqrt{q}}(2)$[11]

$$c^1_{\mu\nu} := \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}_{\mu\nu}, \qquad c^2_{\mu\nu} := \frac{1}{\sqrt{1+q}}\begin{pmatrix} 0 & \sqrt{q} \\ 1 & 0 \end{pmatrix}_{\mu\nu}, \qquad c^3_{\mu\nu} := \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}_{\mu\nu} \tag{14}$$

and $c^i_{\mu\nu} = c^{\mu\nu}_i$ the matrix elements $M^i{}_j$ of $SO_q(3)$ are given in terms of $SU_{\sqrt{q}}(2)$ elements $m^{\mu}{}_{\nu}$ by

$$M^i{}_j := c^i_{\mu\nu} m^{\mu}{}_{\rho} m^{\nu}{}_{\sigma} c^{\rho\sigma}_j. \tag{15}$$

This is verified by checking the $\hat{R}$-matrix of the $SO_q(3)$ group to be :

$$\frac{1}{q}\eta^i{}_{i'}\eta^j{}_{j'}c^i_{\mu\rho}c^j_{\sigma\lambda}\hat{r}^{\mu\nu}{}_{\mu'\nu'}\hat{r}^{\rho\sigma}{}_{\nu\tau}\hat{r}^{\tau\lambda}{}_{\tau'\lambda'}\hat{r}^{\nu'\tau'}{}_{\rho'\sigma'}c^{\mu'\rho'}_k c^{\sigma'\lambda'}_l, \tag{16}$$

where $\hat{r}$ denotes the $\hat{R}$-matrix of the $SU_{\sqrt{q}}(2)$ group and $\eta = \mathrm{diag}(1, i, 1)$, i being the imaginary unit. It just produces a base change to more convenient coordinates. Note as well the decomposition of the symmetric projector $\hat{S}^{\mu\nu}_{\sqrt{q}\rho\sigma} = c^{\mu\nu}_i c^i_{\rho\sigma}$.

In order to clear out nasty indices we want to make use of a graphical notation, which has been given in [10]. With

$$c^i_{\mu\nu} := \tag{17}$$



and the equality



(18)

we can disentangle the matrix:



(19)

This is the $\hat{R}$-matrix of $SO_q(3)$ since obviously

$$C^{ij} = \eta^{i}{}_{i'}\eta^{j}{}_{j'}$$



(20)

and

$$\hat{A}_q{}^{ij}{}_{kl} = \eta^{i}{}_{i'}\eta^{j}{}_{j'}$$



$$\eta^{k'}{}_{k}\eta^{l'}{}_{l}$$

(21)

Of course the same construction holds for the $SO_q(2,1)$ group. Then η has to be chosen as identity matrix.

b) Since we know the q-antisymmetrizer we are able to find the $SO_q(3)$ covariant quantum plane in terms of spinor variables. To obtain sufficiently many degrees of freedom we have to take at least two copies of q-spinors x and y having the same commutation relations with m. This provides an extended inhomogeneous algebra called $ISU^{ex}_{\sqrt{q}}(2)$. We want to mention that the extended algebra does not have a correct coalgebra structure. This is not important for the algebraic embedding. The 3-dimensional quantum space has the form:

$z^i = {\eta^i}_j$



(22)

We fix the x, y- relations such that the element $\varepsilon_{\nu\mu}\, x^\nu y^\mu$ commutes with the coordinate functions and get



(23)

We still have to prove that $\hat{A}_q{}^{ij}{}_{kl} z^i z^j$vanishes. This follows from the equation



$= 0,$ (24)

or



(25)

Next we observe that



(26)

Now we have to take a look to the * - operation. The behaviour is for q complex quite different from q real, since $(c^i_{\mu\nu})^* = c^i_{\nu\mu}$ in the first and $(c^i_{\mu\nu})^* = c^i_{\mu\nu}$ in the second case. From this we see immediately that the algebraic embedding in the case of $ISO_q(2,1)$ respects the * - operation, whereas for the $ISO_q(3)$ quantum group we have to examine the * - structure more closely. At first for the homogeneous part we have:

$$\kappa(c^i_{\mu\nu} m^\mu{}_\rho m^\nu{}_\sigma c^{\rho\sigma}_j) = c^i_{\mu\nu}(m^\sigma{}_\nu)^*(m^\rho{}_\mu)^* c^{\rho\sigma}_j = \left(c^j_{\rho\sigma} m^\rho{}_\mu m^\sigma{}_\nu c^{\mu\nu}_i\right)^* . \tag{27}$$

To use the graphical technique for the translations we have to introduce the "transposed c–matrix "c^T:

$(c^T)_i^{\nu\mu}$ =



= $c_i^{\mu\nu}$ (28)

Of course $(z^i)^* =: \bar{z}_i = (c^T_i)^{\mu\nu}\bar{x}_\mu\bar{x}_\nu$ Calculating now the mixed commutation relations one has:



(29)

To reproduce the correct $z - \bar{z}$ relations which use diagramms similar to that of (29) we have to redefine z as $z \longrightarrow z^i = x^\mu x^\nu c^i_{\mu\nu}\bar{\omega}^n$. Now the calculation is similar to the one above and we find $n = 2/3$. This relation finishes the prove of the algebraic embedding.

Remark:

a) The antipode of the coordinates z just differs by a minus sign, when expressed in terms of $\kappa(x)$.

b) The Coproduct of the coordinates z can't be embedded by obvious reasons.

4 Hilbert space representation for $ISO_q(3)$

The problem to find a representation for $ISO_q(3)$ and $ISO_q(2,1)$ is now reduced to that of representing $ISU^{ex}_{\sqrt{q}}(2)$ and $ISL^{ex}_q(2,R)$.

Here we restrict ourselves to the representation of $ISO_q(3)$. (We want to mention that $SL_q(2,R)$ does not exist on the Hilbert space level anyhow [12].) As well we confine the value of q to $(0,1)$. The case $q>1$ is however isomorphic. The relations of $SU_{\sqrt{q}}(2)$ are $(\mu=\sqrt{q})$:

$$\begin{array}{lll} \alpha\gamma=\mu\gamma\alpha, & \alpha\gamma^*=\mu\gamma^*\alpha \quad , & \gamma\gamma^*=\gamma^*\gamma, \\ \alpha^*\alpha+\gamma^*\gamma=1, & & \alpha\alpha^*+\mu^2\gamma^*\gamma=1 \end{array} \tag{30}$$

The second coordinate has the following commutation relations: (Remember that in our convention the quantum plane coordinates are $x^{(\tau)}, \tau=1,2.$)

$$\begin{array}{rlrl} x^{(2)}\alpha & =\mu^{-1}\alpha x^{(2)}, & x^{(2)}\gamma = & \mu\gamma x^{(2)} \\ x^{(2)}\alpha^* & =\mu\alpha^* x^{(2)}, & x^{(2)}\gamma^* = & \mu^{-1}\gamma^* x^{(2)} \end{array} \tag{31}$$

Applying the antipode to (31) gives

$$\begin{array}{rlrl} \alpha^*\kappa(x^{(2)}) & =\mu^{-1}\kappa(x^{(2)})\alpha^*, & \gamma\kappa(x^{(2)}) = & \mu\kappa(x^{(2)})\gamma \\ \alpha\kappa(x^{(2)}) & =\mu\kappa(x^{(2)})\alpha, & \gamma^*\kappa(x^{(2)}) = & \mu^{-1}\kappa(x^{(2)})\gamma^*. \end{array} \tag{32}$$

Those relations hold for any $SU_{\sqrt{q}}(2)$-covariant plane.

We have other relations for the functions on the quantum planes $x^{(\tau)}$ and $y^{(\tau)}, \tau = 1,2$:

$$\begin{array}{rclrcl} x^{(2)}y^{(2)} & = & \mu y^{(2)}x^{(2)}, & \kappa(x^{(2)})\kappa(y^{(2)}) & = & \mu^{-1}\kappa(y^{(2)})\kappa(x^{(2)}), \\ x^{(\tau)}\kappa(x^{(\delta)}) & = & \mu^{-2}\kappa(x^{(\delta)})x^{(\tau)}, & x^{(\tau)}\kappa(y^{(\delta)}) & = & \mu^{-1}\kappa(y^{(\delta)})x^{(\tau)}, \\ \kappa(x^{(2)})\overline{\kappa(y^{(2)})} & = & \mu^2\overline{\kappa(y_2)}\kappa(x_2), & & & \\ \overline{a_\nu}\kappa(b^{(\tau)}) & = & \kappa(b^{(\tau)})\overline{a_\nu}, & a,b\,\epsilon\,\{x,y\}, & & \\ \multicolumn{6}{c}{y^{(\tau)}\kappa(x^{(\delta)}) = \mu^{-1}\kappa(x^{(\delta)})y^\tau + (\mu^{-2}-1)\kappa(y^{(\delta)})x^{(\tau)},} \end{array} \tag{33}$$

where $\delta,\tau\,\epsilon\,\{1,2\}$.
These and the conjugated relations together with the obvious relations for ω contain the whole algebraic information of $ISU_{\sqrt{q}}(2)$ since:

$$a^{(1)} = (\mu\gamma)^{(-1)}\left(\omega\kappa\left(a^{(2)}\right)+\alpha a^{(2)}\right), \qquad a,b\,\epsilon\,\{x,y\}. \tag{34}$$

Remember that γ is an invertible element.

The algebra may still be simplyfied by a nonlinear transformation in the functions of coordinates. With $Q^4=\mu^2=q$ and $v^3=\omega$ we define:

$$\begin{array}{rclrcl} \rho_1 & = & \bar{v}^2\gamma^{-1}x^{(2)} & \Theta_1 & = & q^{-1}v\gamma^{-1}\kappa(x^{(2)}) \\ \rho_2 & = & \bar{v}^2\gamma^{-1}y^{(2)} & \Theta_2 & = & q^{-1}v\gamma^{-1}\kappa(y^{(2)}) \end{array} \tag{35}$$

All algebraic relations with coordinate-functions are given by:

$$\begin{array}{rclrclrcl}
\rho_i\alpha &=& Q\alpha\rho_i & \Theta_i\alpha &=& Q\alpha\Theta_i & \Theta_i\Theta_i^* &=& \Theta_i^*\Theta_i \\
\rho_i\alpha^* &=& Q^{-1}\alpha^*\rho_i & \Theta_i\alpha^* &=& Q^{-1}\alpha^*\Theta_i & \rho_i\rho_i^* &=& Q^2\rho_i^*\rho_i \\
\rho_i\gamma &=& Q\gamma\rho_i & \Theta_i\gamma &=& Q^{-1}\gamma\Theta_i & \rho_i\Theta_i &=& Q^{-3}\Theta_i\rho_i \\
\rho_i\gamma^* &=& Q^{-1}\gamma^*\rho_i & \Theta_i\gamma^* &=& Q\gamma^*\Theta_i & \rho_i^*\Theta_i &=& Q^{-1}\Theta_i\rho_i^*
\end{array} \tag{36}$$

with $i = 1, 2$.

$$\begin{array}{rclrcl}
\Theta_1\Theta_2 &=& Q^2\Theta_2\Theta_1 & \rho_1\rho_2^* &=& Q^2\rho_2^*\rho_1 \\
\rho_1\rho_2 &=& Q^2\rho_2\rho_1 & \Theta_1\rho_2^* &=& Q\rho_2^*\Theta_1 \\
\rho_1\Theta_2 &=& Q^{-1}\Theta_2\rho_1 & \Theta_2\rho_1^* &=& Q\rho_1^*\Theta_2 \\
\Theta_1\rho_2 &=& (Q^2 - Q^{-2})\Theta_2\rho_1 + Q\rho_2\Theta_1 & \Theta_1\Theta_2^* &=& \Theta_2^*\Theta_1
\end{array} \tag{37}$$

It is easy to find a maximal real subalgebra $\mathcal{D}$ of commuting elements. In order to rely the representation of $ISO_q(3)$ to that of $SU_{\sqrt{q}}(2)$ we choose:

$$\mathcal{D} := \{\alpha\alpha^*, \gamma\gamma^*, \rho_1\rho_1^*, \Theta_1\Theta_1^*, \Theta_2\Theta_2^*, v\}. \tag{38}$$

These operators are used to label the eigenvectors of the Hilbertspace. They are normalised so that the representation is given by:

$$\begin{array}{rcl}
\pi(\alpha)|n,m,k,r,s,v\rangle &=& \sqrt{1-Q^{4n}}|n-1,m,k,r,s,v\rangle \\
\pi(\alpha^*)|n,m,k,r,s,v\rangle &=& \sqrt{1-Q^{4(n+1)}}|n+1,m,k,r,s,v\rangle \\
\pi(\gamma)|n,m,k,r,s,v\rangle &=& Q^{2n}|n,m-1,k,r,s,v\rangle \\
\pi(\gamma^*)|n,m,k,r,s,v\rangle &=& Q^{2n+2}|n,m+1,k,r,s,v\rangle \\
\pi(v)|n,m,k,r,s,v\rangle &=& |n,m,k-1,r,s,v\rangle \\
\pi(v^*)|n,m,k,r,s,v\rangle &=& |n,m,k+1,r,s,v\rangle \\
\pi(\rho_1)|n,m,k,r,s,v\rangle &=& Q^{-(n+m+k)+r+2s-v}|n,m,k,r-1,s,v\rangle \\
\pi(\rho_1^*)|n,m,k,r,s,v\rangle &=& Q^{-(n+m+k)+r+1+2s-v}|n,m,k,r+1,s,v\rangle \\
\pi(\rho_2)|n,m,k,r,s,v\rangle &=& Q^{-(n+m+k)+2r+3s-v}|n,m,k,r-1,s,v-1\rangle \\
\pi(\rho_2^*)|n,m,k,r,s,v\rangle &=& Q^{-(n+m+k)+2r+1+3s-v}|n,m,k,r+1,s,v+1\rangle \\
\pi(\Theta_1)|n,m,k,r,s,v\rangle &=& Q^{-(n-m+k+r+v)}|n,m,k,r,s-1,v\rangle \\
\pi(\Theta_1^*)|n,m,k,r,s,v\rangle &=& Q^{-(n-m+k+r+v)}|n,m,k,r,s+1,v\rangle \\
\pi(\Theta_2)|n,m,k,r,s,v\rangle &=& Q^{-(n-m+k+r-s+v)}|n,m,k,r,s-1,v-1\rangle \\
\pi(\Theta_2^*)|n,m,k,r,s,v\rangle &=& Q^{-(n-m+k+r-s+v)}|n,m,k,r,s+1,v+1\rangle
\end{array} \tag{39}$$

5 Conclusion

We have given the algebraic embedding of two q-euclidean groups in three dimensions We have given an irreducible Hilbert space representation for the function algebra of $ISO_q(3)$. With a reasoning along the lines of [6] it should be possible to prove that $ISO_q(3)$ exists on a C^*-algebra level.

Acknowledgments
We would like to thank R. Engeldinger, J. Seifert and W. Weich for valuable discussions.

References

[1] L.D. Faddeev, N.Yu. Reshetikhin and L.A. Takhtajan, *Algebra and Analysis* **1**, 178 (1987).

[2] M. Schlieker, W. Weich, R. Weixler, "Inhomogeneous Quantum Groups." , *Z. Phys. C* **53**, 79 (1992).

[3] R.O. Weixler, "Inhomogene Quantengruppen", Thesis, Shaker, Aachen (1994).

[4] T. Kobayashi, H. T. Sato, "Quantum Deformation of $igl(n)$ Algebra and Quantum Space", Preprint LMU-TPW 94-27.

[5] B. Drabant, "Braided Bosonization and Inhomogeneous Quantum Groups", Report 94-28, Preprint (1994).

[6] S. L. Woronowicz, "Unbounded Elements Affiliated with C^*-Algebras and Non-Compact Quantum Groups", *Comm. Math. Phys.* **136**, 399 (1991).

[7] G. Fiore, "The Euclidean Hopf algebra $u_q(e_n)$ and its fundamental representation", Preprint, LMU-TP 94-11, (1994).

[8] S. Majid, "Cross Products by Braided Groups and Bosonization", *J.Algebra* **163**, 165 (1994).

[9] U. Carow-Watamura, M. Schlieker, M. Scholl, S. Watamura, "Tensor Representations of the Quantum Group $SL_q(2,\mathbb{C})$ and Quantum Minkowsky Space", *Z. Phys. C* **48**, 159 (1990).

[10] U. Carow-Watamura, M. Schlieker, M. Scholl, S. Watamura, "A Quantum Lorentz Group", *J. Mod. Phys. A*, **6**, 3081 (1991).

[11] W. Weich, "$SO_q(3)$-Symmetric Quantum Mechanics", Preprint LMU-TPW 94-7, (1994).

[12] K. Schmüdgen, " Integrable Operator Representations of $\mathbf{R}_q^2, X_{q,\gamma}$ and $SL_q(2,R)$", *Commun. Math. Phys.* **159**, 217 (1994). .

"Bicovariant Differential Calculus on Quantum Groups $SU_q(N)$ and $SO_q(N)$, *Commun. Math. Phys.* **142**, 605 (1990).